P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Species/Strain: MICE/B6C3F1

Test Type: CHRONIC

F1_M3

NTP Study Number: C20515

Lock Date: 01/12/2011

Cage Range: ALL

Route: RESPIRATORY EXPOSURE WHOLE BODY

Date Range: ALL

Reasons For Removal: ALL

Removal Date Range: ALL

Treatment Groups: Include ALL

Study Gender: Both

TDMSE Version: 3.0.1.1_001

PWG Approval Date: NONE

Date Report Requested: 03/08/2013 **Time Report Requested:** 14:13:47 **First Dose M/F:** 05/05/08 / 05/05/08

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

Angiectasis

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Lab: BNW

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Disposition Summary					
Animals Initially In Study Early Deaths	50	50	50	50	
Moribund Sacrifice	5	12	7	4	
Natural Death	6	7	7	6	
Survivors					
Terminal Sacrifice	39	31	36	40	
Animals Examined Microscopically	50	50	50	50	
ALIMENTARY SYSTEM					
Esophagus	(50)	(50)	(50)	(50)	
Gallbladder	(44)	(44)	(44)	(44)	
Cytoplasmic Alteration	. ,		. ,	1 (2%)	
Inflammation, Suppurative	1 (2%)			1 (2%)	
Intestine Large, Cecum	(50)	(50)	(49)	(49)	
Hemorrhage			1 (2%)		
Intestine Large, Colon	(50)	(50)	(50)	(50)	
Ulcer		1 (2%)			
Intestine Large, Rectum	(50)	(50)	(50)	(49)	
Cyst		1 (2%)			
Fibrosis		1 (2%)			
Intestine Small, Duodenum	(49)	(48)	(48)	(47)	
Artery, Infiltration Cellular, Lymphocyte	1 (2%)				
Intestine Small, Ileum	(50)	(50)	(49)	(48)	
Hyperplasia, Lymphoid				1 (2%)	
Inflammation, Chronic Active			2 (4%)	2 (4%)	
Epithelium, Hyperplasia	1 (2%)				
Lymphoid Follicle, Hyperplasia	1 (2%)				
Peyer's Patch, Hyperplasia, Lymphoid	1 (2%)				
Intestine Small, Jejunum	(50)	(48)	(49)	(48)	
Inflammation, Chronic Active			1 (2%)		
Serosa, Inflammation, Granulomatous				1 (2%)	
Liver	(50)	(50)	(50)	(49)	

a - Number of animals examined microscopically at site and number of animals with lesion

1 (2%)

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Basophilic Focus	1 (2%)	3 (6%)		4 (8%)	
Clear Cell Focus	10 (20%)	9 (18%)	3 (6%)	6 (12%)	
Congestion		1 (2%)	1 (2%)		
Cyst		1 (2%)	1 (2%)		
Eosinophilic Focus	14 (28%)	3 (6%)	20 (40%)	17 (35%)	
Fatty Change	24 (48%)	22 (44%)	24 (48%)	25 (51%)	
Hepatodiaphragmatic Nodule	1 (2%)		1 (2%)		
Hyperplasia	1 (2%)	1 (2%)	1 (2%)		
Infiltration Cellular, Lymphocyte		` ,	1 (2%)		
Inflammation, Chronic	1 (2%)		,		
Inflammation, Chronic Active	,			1 (2%)	
Mixed Cell Focus	6 (12%)	4 (8%)	11 (22%)	5 (10%)	
Necrosis	5 (10%)	7 (14%)	6 (12%)	7 (14%)	
Tension Lipidosis	3 (6%)	2 (4%)	5 (10%)	7 (14%)	
Mesentery	(4)	(5)	(3)	(4)	
Inflammation, Chronic Active	()	1 (20%)	1 (33%)	()	
Arteriole, Inflammation, Chronic Active		1 (20%)	()		
Artery, Inflammation, Chronic Active		,		1 (25%)	
Fat, Necrosis	1 (25%)	2 (40%)	2 (67%)	3 (75%)	
Pancreas	(50)	(50)	(50)	(50)	
Fibrosis	1 (2%)	()	()	()	
Lipomatosis	(/	1 (2%)			
Necrosis		()		1 (2%)	
Acinus, Hypertrophy	1 (2%)	1 (2%)		(= / -)	
Artery, Inflammation, Chronic Active	(/	1 (2%)			
Salivary Glands	(50)	(50)	(50)	(50)	
Infiltration Cellular, Lymphocyte	(5.5)	1 (2%)	()	()	
Stomach, Forestomach	(50)	(50)	(50)	(50)	
Cyst, Squamous	(5.5)	1 (2%)	()	()	
Hyperplasia, Squamous	2 (4%)	2 (4%)	1 (2%)	3 (6%)	
Inflammation, Chronic Active	1 (2%)	2 (4%)	1 (2%)	- (5/5)	
Ulcer	(=/0)	_ (· / • /	1 (2%)		
Stomach, Glandular	(50)	(50)	(50)	(50)	
Erosion	1 (2%)	(00)	(00)	(33)	
Inflammation, Chronic Active	1 (2%)			2 (4%)	
Ulcer	. (=/0)			1 (2%)	

a - Number of animals examined microscopically at site and number of animals with lesion

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Arteriole, Serosa, Inflammation, Chronic Active		1 (2%)			
Glands, Mineralization			2 (4%)	1 (2%)	
Tooth	(50)	(47)	(49)	(50)	
Dysplasia	43 (86%)	35 (74%)	41 (84%)	40 (80%)	
Inflammation, Chronic Active	3 (6%)	1 (2%)	6 (12%)	5 (10%)	
CARDIOVASCULAR SYSTEM					
Blood Vessel	(45)	(49)	(48)	(48)	
Inflammation, Chronic Active	1 (2%)	,	` '	,	
Aorta, Inflammation, Chronic Active	, ,	1 (2%)			
Heart	(50)	(50)	(50)	(50)	
Cardiomyopathy	2 (4%)	3 (6%)	1 (2%)	2 (4%)	
Inflammation, Suppurative			1 (2%)		
Inflammation, Chronic Active				1 (2%)	
Thrombosis	1 (2%)			2 (4%)	
Artery, Inflammation, Chronic Active		1 (2%)		1 (2%)	
Valve, Fibrosis			1 (2%)		
ENDOCRINE SYSTEM					
Adrenal Cortex	(49)	(50)	(50)	(50)	
Hyperplasia	24 (49%)	23 (46%)	26 (52%)	24 (48%)	
Capsule, Inflammation, Chronic Active	1 (2%)				
Adrenal Medulla	(48)	(50)	(50)	(50)	
Hyperplasia		1 (2%)		2 (4%)	
Islets, Pancreatic	(50)	(50)	(50)	(50)	
Parathyroid Gland	(34)	(28)	(27)	(21)	
Cyst	1 (3%)				
Pituitary Gland	(50)	(50)	(50)	(48)	
Angiectasis				1 (2%)	
Pars Distalis, Cyst	1 (2%)	- 4		1 (2%)	
Pars Distalis, Hyperplasia	5 (10%)	8 (16%)	7 (14%)	2 (4%)	
Pars Intermedia, Hypertrophy	(50)	1 (2%)	(40)	(50)	
Thyroid Gland	(50)	(50)	(49)	(50)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800) CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Inflammation, Chronic Active			1 (2%)		
Follicular Cell, Hyperplasia			1 (2%)	1 (2%)	
GENERAL BODY SYSTEM					
Tissue NOS	(0)	(2)	(0)	(1)	
GENITAL SYSTEM					
Epididymis	(50)	(50)	(50)	(50)	
Inflammation, Chronic Active	1 (2%)	1 (2%)		2 (4%)	
Spermatocele	(0)	1 (2%)	(0)	(4)	
Penis	(0)	(0)	(0)	(1)	
Congestion	(50)	(40)	(40)	1 (100%)	
Preputial Gland Ectasia	(50)	(49)	(48)	(49)	
	8 (16%)	4 (8%)	7 (15%)	7 (14%)	
Inflammation, Chronic Active	9 (18%)	4 (8%)	11 (23%)	4 (8%)	
Prostate Seminal Vesicle	(50)	(49)	(50) (50)	(50) (50)	
Dilatation	(50)	(50)	(50) 1 (2%)		
Testes	(50)	(50)	(50)	1 (2%) (50)	
Angiectasis	(30)	(50)	(50)	1 (2%)	
Nuclear Alteration, Focal		1 (2%)		1 (270)	
Germinal Epithelium, Atrophy	1 (2%)	4 (8%)		3 (6%)	
	. (=76)	. (0,76)		(6/3)	
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(50)	(50)	(50)	
Hyperplasia			1 (2%)		
Hyperplasia, Granulocytes				1 (2%)	
Lymph Node	(4)	(3)	(1)	(1)	
Lumbar, Infiltration Cellular, Mixed Cell	1 (25%)				
Lumbar, Inflammation, Pyogranulomatous				1 (100%)	
Lumbar, Lymphoid Follicle, Hyperplasia	1 (25%)				

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Lymphoid Follicle, Renal, Hyperplasia	1 (25%)				
Pancreatic, Inflammation, Pyogranulomatous				1 (100%)	
Renal, Inflammation, Pyogranulomatous				1 (100%)	
Lymph Node, Bronchial	(26)	(31)	(26)	(18)	
Infiltration Cellular, Polymorphonuclear	1 (4%)				
Lymph Node, Mandibular	(31)	(36)	(34)	(33)	
Hyperplasia, Lymphoid		2 (6%)	1 (3%)		
Infiltration Cellular, Plasma Cell		1 (3%)			
Inflammation, Pyogranulomatous				1 (3%)	
Lymph Node, Mediastinal	(39)	(39)	(32)	(35)	
Hematopoietic Cell Proliferation			1 (3%)		
Hyperplasia, Plasma Cell			1 (3%)		
Infiltration Cellular, Mixed Cell			1 (3%)		
Inflammation, Pyogranulomatous				1 (3%)	
Lymphoid Follicle, Hyperplasia	1 (3%)			,	
Lymph Node, Mesenteric	(49)	(50)	(48)	(49)	
Fibrosis	1 (2%)	,	, ,	, ,	
Hemorrhage	, ,	1 (2%)	1 (2%)		
Hyperplasia, Plasma Cell			1 (2%)		
Infiltration Cellular, Histiocyte			1 (2%)		
Inflammation, Pyogranulomatous			, ,	1 (2%)	
Lymphoid Follicle, Hyperplasia	1 (2%)			,	
Spleen	(50)	(50)	(49)	(49)	
Congestion	,	,	1 (2%)	` ,	
Fibrosis		1 (2%)	,		
Hematopoietic Cell Proliferation	1 (2%)	1 (2%)	3 (6%)	1 (2%)	
Lymphoid Follicle, Hyperplasia	3 (6%)	2 (4%)	1 (2%)	1 (2%)	
Thymus	(41)	(38)	(39)	(41)	
Atrophy	. ,	. ,	1 (3%)	1 (2%)	
INTEGUMENTARY SYSTEM					
Mammary Gland	(1)	(0)	(1)	(0)	
Skin	(50)	(49)	(50)	(50)	
Abscess				1 (2%)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Angiectasis			1 (2%)		
Inflammation, Chronic Active	2 (4%)	4 (8%)	1 (2%)	3 (6%)	
Ulcer	1 (2%)	4 (8%)	2 (4%)		
Epidermis, Hyperplasia				1 (2%)	
Sebaceous Gland, Hyperplasia		1 (2%)			
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Fibro-Osseous Lesion	7 (14%)	7 (14%)	10 (20%)	7 (14%)	
Cranium, Fracture		1 (2%)			
Cranium, Inflammation, Chronic Active	3 (6%)	2 (4%)	3 (6%)	5 (10%)	
Joint, Vertebra, Fracture		1 (2%)			
Maxilla, Necrosis		1 (2%)			
Skeletal Muscle	(1)	(3)	(2)	(3)	
Infiltration Cellular, Lymphocyte		1 (33%)			
Infiltration Cellular, Mixed Cell				1 (33%)	
Inflammation, Chronic Active				1 (33%)	
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Inflammation, Suppurative	(,	1 (2%)	,	
Thrombosis			1 (2%)		
Arteriole, Inflammation, Chronic Active		1 (2%)	, ,		
Ventricle, Dilatation		` ,	1 (2%)		
Venule, Infiltration Cellular, Lymphoid		1 (2%)	, ,		
Peripheral Nerve	(1)	(2)	(1)	(2)	
Spinal Cord	(1)	(3)	(1)	(2)	
RESPIRATORY SYSTEM					
Larynx	(50)	(50)	(49)	(50)	
Hyperplasia	(55)	(30)	1 (2%)	(/	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Lab: BNW

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3
Inflammation, Suppurative		1 (2%)		1 (2%)
Inflammation, Chronic Active		2 (4%)	3 (6%)	8 (16%)
Metaplasia, Squamous		50 (100%)	49 (100%)	50 (100%)
Necrosis		1 (2%)	1 (2%)	1 (2%)
Lung	(50)	(50)	(50)	(50)
Congestion			1 (2%)	
Erythrophagocytosis		1 (2%)		
Hemorrhage			2 (4%)	1 (2%)
Infiltration Cellular, Histiocyte	5 (10%)	5 (10%)	1 (2%)	9 (18%)
Inflammation, Acute			1 (2%)	
Thrombosis	1 (2%)			
Alveolar Epithelium, Hyperplasia	4 (8%)	4 (8%)	6 (12%)	7 (14%)
Bronchiole, Hyperplasia	11 (22%)	11 (22%)	32 (64%)	44 (88%)
Interstitium, Inflammation, Chronic Active	1 (2%)		1 (2%)	1 (2%)
Perivascular, Inflammation, Chronic Active	1 (2%)	1 (2%)	1 (2%)	1 (2%)
Nose	(50)	(50)	(50)	(50)
Inflammation, Chronic Active	6 (12%)	8 (16%)	4 (8%)	12 (24%)
Necrosis	2 (4%)	2 (4%)		1 (2%)
Nasopharyngeal Duct, Necrosis				1 (2%)
Olfactory Epithelium, Accumulation, Hyaline Droplet	4 (8%)	31 (62%)	43 (86%)	49 (98%)
Olfactory Epithelium, Atrophy	1 (2%)			4 (8%)
Olfactory Epithelium, Metaplasia, Respiratory	7 (14%)	15 (30%)	25 (50%)	37 (74%)
Olfactory Epithelium, Necrosis		1 (2%)		
Respiratory Epithelium, Accumulation, Hyaline Droplet	7 (14%)	36 (72%)	50 (100%)	50 (100%)
Respiratory Epithelium, Hyperplasia	38 (76%)	44 (88%)	36 (72%)	41 (82%)
Respiratory Epithelium, Metaplasia, Squamous		3 (6%)		1 (2%)
Turbinate, Atrophy				1 (2%)
Turbinate, Degeneration	1 (2%)	1 (2%)		
Pleura	(1)	(0)	(0)	(0)
Trachea	(50)	(50)	(47)	(50)
Ulcer				1 (2%)

SPECIAL SENSES SYSTEM

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE MALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Eye	(50)	(50)	(50)	(50)	
Atrophy	1 (2%)	,	,	,	
Cornea, Hyperplasia, Squamous				1 (2%)	
Cornea, Inflammation, Chronic Active		4 (8%)	2 (4%)	2 (4%)	
Cornea, Vacuolization Cytoplasmic	2 (4%)	` ,	, ,	,	
Lens, Degeneration	1 (2%)	1 (2%)		1 (2%)	
Harderian Gland	(50)	(50)	(49)	(50)	
Cyst	,	1 (2%)	,	,	
Hyperplasia	2 (4%)	1 (2%)	3 (6%)	2 (4%)	
URINARY SYSTEM					
Kidney	(50)	(50)	(50)	(50)	
Congestion	1 (2%)				
Cyst	4 (8%)	2 (4%)	3 (6%)	5 (10%)	
Dilatation	1 (2%)	` ,	, ,	, ,	
Infarct	3 (6%)	1 (2%)	4 (8%)	2 (4%)	
Inflammation, Suppurative			1 (2%)		
Inflammation, Chronic Active	1 (2%)				
Metaplasia, Osseous	1 (2%)	1 (2%)			
Nephropathy	48 (96%)	48 (96%)	45 (90%)	48 (96%)	
Artery, Hemorrhage	1 (2%)				
Artery, Inflammation, Chronic Active	1 (2%)	3 (6%)			
Capsule, Fibrosis		1 (2%)			
Capsule, Hemorrhage		1 (2%)			
Papilla, Necrosis		1 (2%)			
Pelvis, Inflammation, Suppurative		1 (2%)			
Ureter	(0)	(0)	(0)	(1)	
Inflammation, Chronic Active				1 (100%)	
Urinary Bladder	(50)	(50)	(50)	(49)	

^{***} END OF MALE ***

a - Number of animals examined microscopically at site and number of animals with lesion

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

Hyperplasia

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800) CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Lab: BNW

6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Disposition Summary					
Animals Initially In Study	50	50	50	50	
Early Deaths					
Accidentally Killed				1	
Moribund Sacrifice	7	8	10	11	
Natural Death	4	5	3	5	
Survivors					
Terminal Sacrifice	39	37	37	33	
Animals Examined Microscopically	50	50	50	50	
LIMENTARY SYSTEM					
Esophagus	(50)	(50)	(50)	(50)	
Gallbladder	(46)	(49)	(41)	(41)	
Epithelium, Hyperplasia		1 (2%)			
Intestine Large, Cecum	(50)	(50)	(49)	(50)	
Hyperplasia		1 (2%)			
Inflammation, Chronic Active			1 (2%)		
Intestine Large, Colon	(49)	(50)	(49)	(50)	
Intestine Large, Rectum	(50)	(50)	(50)	(50)	
Inflammation, Chronic Active			1 (2%)	1 (2%)	
Intestine Small, Duodenum	(47)	(50)	(50)	(49)	
Intestine Small, Ileum	(49)	(50)	(49)	(49)	
Inflammation, Chronic Active			1 (2%)		
Intestine Small, Jejunum	(48)	(50)	(48)	(48)	
Liver	(50)	(50)	(50)	(50)	
Amyloid Deposition				1 (2%)	
Angiectasis		2 (4%)	1 (2%)	2 (4%)	
Basophilic Focus	6 (12%)	5 (10%)	3 (6%)	4 (8%)	
Clear Cell Focus	1 (2%)				
Cyst		1 (2%)		1 (2%)	
Degeneration		1 (2%)			
Eosinophilic Focus	6 (12%)	1 (2%)	5 (10%)	13 (26%)	
Fatty Change	3 (6%)	2 (4%)	1 (2%)	5 (10%)	

a - Number of animals examined microscopically at site and number of animals with lesion

1 (2%)

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Infiltration Cellular, Lymphocyte		1 (2%)		1 (2%)	
Infiltration Cellular, Mixed Cell		1 (2%)		, ,	
Inflammation, Chronic Active		` ,	1 (2%)		
Mixed Cell Focus	2 (4%)	4 (8%)	1 (2%)		
Necrosis	2 (4%)		2 (4%)	4 (8%)	
Tension Lipidosis	6 (12%)	5 (10%)	5 (10%)	6 (12%)	
Vacuolization Cytoplasmic	1 (2%)				
Centrilobular, Degeneration, Acute		1 (2%)			
Centrilobular, Hypertrophy			1 (2%)		
Centrilobular, Necrosis		1 (2%)			
Mesentery	(16)	(23)	(21)	(15)	
Angiectasis	1 (6%)	, ,	, ,	,	
Cyst	1 (6%)				
Fibrosis	, ,		1 (5%)		
Inflammation, Suppurative			1 (5%)		
Inflammation, Chronic Active	1 (6%)	1 (4%)	1 (5%)		
Necrosis		1 (4%)	1 (5%)		
Arteriole, Inflammation, Chronic Active			1 (5%)		
Fat, Cyst			1 (5%)		
Fat, Necrosis	11 (69%)	17 (74%)	7 (33%)	8 (53%)	
Vein, Inflammation, Chronic Active	, ,	` ,	, ,	1 (7%)	
Pancreas	(50)	(50)	(50)	(50)	
Cyst		1 (2%)			
Fibrosis	1 (2%)				
Inflammation, Chronic Active	1 (2%)			1 (2%)	
Lipomatosis		1 (2%)		1 (2%)	
Acinus, Atrophy	1 (2%)		1 (2%)	2 (4%)	
Acinus, Hyperplasia				1 (2%)	
Acinus, Hypertrophy			1 (2%)	1 (2%)	
Salivary Glands	(50)	(50)	(50)	(49)	
Stomach, Forestomach	(50)	(50)	(50)	(50)	
Fibrosis	1 (2%)	, ,	, ,	,	
Hyperplasia, Squamous	1 (2%)	1 (2%)	2 (4%)	2 (4%)	
Inflammation, Chronic Active	, ,	, ,	, ,	1 (2%)	
Ulcer		1 (2%)	1 (2%)	1 (2%)	
Arteriole, Inflammation, Chronic Active			1 (2%)	. ,	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Lab: BNW

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Stomach, Glandular	(50)	(50)	(50)	(50)	
Degeneration, Hyaline	1 (2%)				
Fibrosis			1 (2%)		
Inflammation, Chronic Active	1 (2%)				
Ulcer				1 (2%)	
Arteriole, Inflammation, Chronic Active			1 (2%)		
Glands, Mineralization			2 (4%)	1 (2%)	
Tooth	(1)	(0)	(2)	(4)	
Dysplasia	1 (100%)		2 (100%)	3 (75%)	
Inflammation, Chronic Active				2 (50%)	
CARDIOVASCULAR SYSTEM					
Blood Vessel	(46)	(44)	(49)	(44)	
Inflammation, Chronic Active	,	,	1 (2%)	,	
Aorta, Embolus Bacterial	1 (2%)		,	1 (2%)	
Aorta, Inflammation, Suppurative	,			1 (2%)	
Heart	(50)	(50)	(50)	(50)	
Cardiomyopathy	1 (2%)	3 (6%)	8 (16%)	5 (10%)	
Congestion	,	1 (2%)	,	1 (2%)	
Mineralization		,	1 (2%)	` ,	
Necrosis	1 (2%)	1 (2%)	,	1 (2%)	
Thrombosis	,	1 (2%)		, ,	
Artery, Inflammation, Chronic Active	1 (2%)	,			
Capillary, Hyperplasia	1 (2%)				
ENDOCRINE SYSTEM					
Adrenal Cortex	(49)	(50)	(50)	(50)	
Angiectasis	1 (2%)	(30)	(30)	(/	
Atrophy	. (=,0)		1 (2%)	1 (2%)	
Degeneration, Cystic		1 (2%)	. (= / = /	. (=,0)	
Hyperplasia	10 (20%)	5 (10%)	10 (20%)	6 (12%)	
Vacuolization Cytoplasmic	2 (4%)	2 (4%)	. 5 (2070)	2 (4%)	
Adrenal Medulla	(49)	(50)	(49)	(50)	

a - Number of animals examined microscopically at site and number of animals with lesion

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Hyperplasia	2 (4%)	1 (2%)	2 (4%)	4 (8%)	
Islets, Pancreatic	(50)	(50)	(50)	(49)	
Parathyroid Gland	(26)	(29)	(22)	(25)	
Cyst		1 (3%)			
Hyperplasia	1 (4%)				
Pituitary Gland	(50)	(50)	(49)	(50)	
Angiectasis	2 (4%)	1 (2%)		4 (8%)	
Pars Distalis, Cyst		1 (2%)		1 (2%)	
Pars Distalis, Hyperplasia	17 (34%)	21 (42%)	19 (39%)	24 (48%)	
Pars Intermedia, Pars Nervosa, Cyst	, ,	, ,	, ,	1 (2%)	
Thyroid Gland	(50)	(48)	(50)	(50)	
Inflammation, Chronic Active	, ,	1 (2%)	2 (4%)	2 (4%)	
C-cell, Hyperplasia	1 (2%)	1 (2%)	, ,	, ,	
Follicle, Cyst	, ,	1 (2%)			
Follicular Cell, Hyperplasia	1 (2%)	1 (2%)	2 (4%)	3 (6%)	
GENERAL BODY SYSTEM					
Tissue NOS	(3)	(2)	(0)	(0)	
Mediastinum, Infiltration Cellular, Lymphocyte	1 (33%)	(-)	(0)	(5)	
Mediastinum, Thrombosis	1 (33%)				
GENITAL SYSTEM					
Clitoral Gland	(44)	(45)	(43)	(44)	
Hyperplasia, Basal Cell	(/	(15)	(15)	1 (2%)	
Inflammation, Chronic Active	1 (2%)	1 (2%)	1 (2%)	(=72)	
Ovary	(50)	(50)	(48)	(50)	
Angiectasis	3 (6%)	2 (4%)	(10)	3 (6%)	
Cyst	14 (28%)	12 (24%)	12 (25%)	16 (32%)	
Hemorrhage	12 (24%)	16 (32%)	7 (15%)	15 (30%)	
Infiltration Cellular, Lymphocyte	(- · · ·)	1 (2%)	1 (2%)	(,-,	
Inflammation, Chronic Active	1 (2%)	2 (4%)	3 (6%)	2 (4%)	
Mineralization	1 (2%)	- (· / • /	- (0,0)	1 (2%)	
Necrosis	1 (2%)			- (-,-)	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Thrombosis				1 (2%)	
Corpus Luteum, Hyperplasia	3 (6%)	1 (2%)			
Uterus	(50)	(50)	(50)	(50)	
Angiectasis	5 (10%)	3 (6%)	1 (2%)	1 (2%)	
Atrophy	1 (2%)				
Decidual Reaction	1 (2%)				
Dilatation	6 (12%)	5 (10%)	8 (16%)	6 (12%)	
Hemorrhage				1 (2%)	
Inflammation, Suppurative	1 (2%)				
Inflammation, Chronic Active	2 (4%)	2 (4%)	3 (6%)	4 (8%)	
Thrombosis	1 (2%)	•	·		
Endometrium, Hyperplasia, Cystic	28 (56%)	29 (58%)	33 (66%)	34 (68%)	
HEMATOPOIETIC SYSTEM					
Bone Marrow	(50)	(50)	(50)	(50)	
Lymph Node	(11)	(12)	(16)	(10)	
Ectasia				1 (10%)	
Hemorrhage	1 (9%)				
Iliac, Angiectasis		1 (8%)			
Iliac, Ectasia	1 (9%)				
Iliac, Hyperplasia, Lymphoid	1 (9%)	1 (8%)	1 (6%)		
Iliac, Hyperplasia, Plasma Cell				1 (10%)	
Lumbar, Ectasia	1 (9%)	1 (8%)	2 (13%)		
Lumbar, Hyperplasia, Lymphoid		1 (8%)		1 (10%)	
Lumbar, Infiltration Cellular, Histiocyte		1 (8%)			
Lumbar, Infiltration Cellular, Mixed Cell			1 (6%)		
Pancreatic, Inflammation, Chronic Active	1 (9%)		. ,		
Renal, Angiectasis		1 (8%)			
Renal, Ectasia	1 (9%)	, ,	1 (6%)		
Renal, Hyperplasia, Lymphoid	, ,	2 (17%)	` ,		
Renal, Necrosis		, ,	1 (6%)		
Lymph Node, Bronchial	(26)	(27)	(28)	(27)	
Infiltration Cellular, Histiocyte	, ,	,	,	1 (4%)	
Necrosis			1 (4%)	,	

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Filst Dose i

Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Date Report Requested: 03/08/2013

Lymph Node, Mandibular			30 mg/m3	100 mg/m3	
Lymph Node, Mandibulai	(41)	(42)	(43)	(35)	
Necrosis			1 (2%)		
Lymph Node, Mediastinal	(43)	(46)	(40)	(47)	
Hematopoietic Cell Proliferation		1 (2%)			
Hyperplasia, Lymphoid		1 (2%)		2 (4%)	
Infiltration Cellular, Mixed Cell				1 (2%)	
Lymph Node, Mesenteric	(48)	(50)	(49)	(49)	
Angiectasis		1 (2%)	2 (4%)	1 (2%)	
Ectasia			2 (4%)		
Hematopoietic Cell Proliferation		1 (2%)			
Hemorrhage		2 (4%)			
Hyperplasia, Lymphoid			1 (2%)		
Infiltration Cellular, Histiocyte	1 (2%)				
Spleen	(50)	(50)	(50)	(49)	
Hematopoietic Cell Proliferation	4 (8%)	3 (6%)	3 (6%)	2 (4%)	
Hyperplasia, Lymphoid	, ,	1 (2%)	1 (2%)	1 (2%)	
Infiltration Cellular, Mast Cell				1 (2%)	
Lymphoid Follicle, Hyperplasia			1 (2%)	, ,	
Thymus	(44)	(46)	(45)	(45)	
Angiectasis	, ,		1 (2%)	. ,	
Atrophy	2 (5%)		, ,	1 (2%)	
Hyperplasia, Lymphoid	1 (2%)			, ,	
Necrosis				1 (2%)	
INTEGUMENTARY SYSTEM					
Mammary Gland	(50)	(50)	(50)	(50)	
Hyperplasia	2 (4%)			1 (2%)	
Skin	(50)	(50)	(50)	(50)	
Infiltration Cellular, Mast Cell				1 (2%)	
Inflammation, Chronic Active			1 (2%)	2 (4%)	
Ulcer	1 (2%)		1 (2%)	3 (6%)	
Epidermis, Hyperplasia	1 (2%)				
Subcutaneous Tissue, Fibrosis	·		1 (2%)		

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
MUSCULOSKELETAL SYSTEM					
Bone	(50)	(50)	(50)	(50)	
Fibro-Osseous Lesion	23 (46%)	25 (50%)	27 (54%)	23 (46%)	
Cartilage, Femur, Metaphysis, Hyperplasia Cranium, Inflammation, Chronic Active Femur, Fracture	1 (2%) 1 (2%) 1 (2%)		2 (4%)	1 (2%)	
Femur, Tibia, Arthrosis	()			1 (2%)	
Maxilla, Hyperostosis			1 (2%)		
Pelvis, Degeneration				1 (2%)	
Vertebra, Fracture				1 (2%)	
Skeletal Muscle	(3)	(3)	(1)	(4)	
Degeneration				1 (25%)	
Infiltration Cellular, Lymphocyte	3 (100%)	1 (33%)			
Infiltration Cellular, Polymorphonuclear				1 (25%)	
Arteriole, Inflammation, Chronic Active			1 (100%)		
NERVOUS SYSTEM					
Brain	(50)	(50)	(50)	(50)	
Compression	3 (6%)	1 (2%)	2 (4%)	3 (6%)	
Meninges, Inflammation, Chronic Active	,	,	,	1 (2%)	
Peripheral Nerve	(2)	(3)	(1)	(3)	
Degeneration	,	1 (33%)	,	1 (33%)	
Infiltration Cellular, Mixed Cell	1 (50%)	, ,		,	
Infiltration Cellular, Polymorphonuclear	, ,	1 (33%)			
Spinal Cord	(3)	(3)	(1)	(3)	
Infiltration Cellular, Lymphocyte		1 (33%)	,	,	
RESPIRATORY SYSTEM					
Larynx Inflammation, Chronic Active	(49)	(49)	(50)	(50) 10 (20%)	
				()	

a - Number of animals examined microscopically at site and number of animals with lesion

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

Test Type: CHRONIC

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Necrosis				4 (8%)	
Lung	(50)	(50)	(50)	(50)	
Congestion				1 (2%)	
Hemorrhage	1 (2%)	1 (2%)	1 (2%)	2 (4%)	
Infiltration Cellular, Histiocyte	4 (8%)	1 (2%)	1 (2%)	6 (12%)	
Infiltration Cellular, Lymphocyte		1 (2%)			
Infiltration Cellular, Mixed Cell				1 (2%)	
Metaplasia, Osseous		1 (2%)			
Thrombosis				1 (2%)	
Alveolar Epithelium, Hyperplasia	4 (8%)	2 (4%)	5 (10%)	4 (8%)	
Alveolus, Infiltration Cellular, Histiocyte	1 (2%)				
Bronchiole, Hyperplasia	7 (14%)	4 (8%)	22 (44%)	41 (82%)	
Interstitium, Inflammation, Chronic Active	1 (2%)			1 (2%)	
Perivascular, Inflammation, Chronic Active	4 (8%)	3 (6%)		11 (22%)	
Nose	(50)	(49)	(50)	(50)	
Inflammation, Chronic Active	9 (18%)		3 (6%)	5 (10%)	
Necrosis	1 (2%)	1 (2%)	2 (4%)	1 (2%)	
Olfactory Epithelium, Accumulation, Hyaline Droplet	25 (50%)	40 (82%)	50 (100%)	49 (98%)	
Olfactory Epithelium, Atrophy	1 (2%)	1 (2%)		1 (2%)	
Olfactory Epithelium, Metaplasia, Respiratory	3 (6%)	4 (8%)	12 (24%)	23 (46%)	
Respiratory Epithelium, Accumulation, Hyaline Droplet	34 (68%)	48 (98%)	50 (100%)	50 (100%)	
Respiratory Epithelium, Hyperplasia	39 (78%)	48 (98%)	39 (78%)	39 (78%)	
Respiratory Epithelium, Metaplasia, Squamous		1 (2%)		1 (2%)	
Respiratory Epithelium, Necrosis		1 (2%)			
Trachea	(49)	(49)	(50)	(50)	
SPECIAL SENSES SYSTEM					
Eye	(50)	(50)	(50)	(50)	
Atrophy			2 (4%)		
Cornea, Inflammation, Chronic Active		1 (2%)		1 (2%)	
Lens, Degeneration	2 (4%)		2 (4%)	1 (2%)	
Harderian Gland	(50)	(50)	(50)	(50)	
Atrophy	1 (2%)		1 (2%)		

a - Number of animals examined microscopically at site and number of animals with lesion

Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Metal working fluids (CIMSTAR 3800)

CAS Number: CIMSTAR3800

Date Report Requested: 03/08/2013 Time Report Requested: 14:13:47 First Dose M/F: 05/05/08 / 05/05/08

Lab: BNW

B6C3F1 MICE FEMALE	Control	10 mg/m3	30 mg/m3	100 mg/m3	
Hyperplasia	2 (4%)	1 (2%)	6 (12%)		
Inflammation, Suppurative	,	,	1 (2%)		
Inflammation, Chronic Active		1 (2%)	1 (2%)		
URINARY SYSTEM					
Kidney	(50)	(50)	(50)	(50)	
Accumulation, Hyaline Droplet	1 (2%)	1 (2%)			
Infarct	1 (2%)	4 (8%)	4 (8%)	1 (2%)	
Inflammation, Suppurative				1 (2%)	
Metaplasia, Osseous		1 (2%)	2 (4%)	1 (2%)	
Necrosis	1 (2%)				
Nephropathy	47 (94%)	46 (92%)	49 (98%)	47 (94%)	
Glomerulus, Amyloid Deposition	1 (2%)			1 (2%)	
Ureter	(0)	(0)	(1)	(0)	
Urethra	(0)	(1)	(0)	(0)	
Urinary Bladder	(50)	(50)	(50)	(50)	
Angiectasis	1 (2%)		1 (2%)		
Arteriole, Inflammation, Chronic Active			1 (2%)		
Artery, Serosa, Inflammation, Chronic Active		1 (2%)	. ,		

*** END OF REPORT ***